



# Technical Data Sheet: hTERT-immortalized White Preadipocytes

<b>ATCC® Number</b>	CRL-4063™
<b>Organism</b>	<i>Homo sapiens</i>
<b>Tissue/Disease Source</b>	Subcutaneous abdominal adipose tissue
<b>Product Description</b>	hTERT immortalized white preadipocytes were isolated from subcutaneous abdominal adipose tissue from a donor with von Hippel-Lindau syndrome.
<b>Application</b>	This hTERT-immortalized primary cell has applications as an in vitro cell model for toxicity studies and the study of obesity and related diseases.




**Figure 1: Cell morphology of Immortalized White Preadipocytes.** Cells were maintained in ATCC recommended culture conditions. High and low confluence images of plated adherent white preadipocytes were taken using Nikon microscope at 10x. Scale bar represents 100 microns.

## Preadipocyte Marker Analysis



**Figure 2: Marker expression by flow cytometry.** Cells were stained for CD90 and CD166 which are common markers for cells of mesenchymal stem cell origin, and CD34, a marker for hematopoietic stem cells. As expected, white preadipocytes were positive for CD90, CD166, but negative for CD34.

## Differentiation of White Preadipocytes into Mature Adipocytes



**Figure 3: Differentiation of hTERT White Preadipocytes from day 0 through day 25.** Images show the morphology of differentiating cells at different timepoints.

### Protocol:

- Culture cells to reach around 70% confluent, seed cells into 6 well plates at a seeding density of 100k cells/cm<sup>2</sup>.
- Observe cells under a phase contrast microscope daily until cells are 100% confluent. Wait 48 additional hours to begin differentiation/induction.
- Replace the cell culture media with Induction Media 1. Incubate the cells for 7 days.
- Switch to Induction Media 2. Grow the cells for additional 14-18 days until Oil droplets can be seen clearly in the mature adipocytes.

### Media Formulations:

#### Induction Media 1:

- Advanced DMEM
- 2% Sigma FBS
- 1% Pen/Strep
- 0.5 uM Human Insulin
- 0.1uM Dexamethasone
- 0.5 mM IBMX
- 33 uM Biotin
- 2 nM T3
- 30 uM Indomethacin
- 17 uM Pantothenate
- 2 uM Rosiglitazone

#### Induction Media 2:

- Advanced DMEM
- 2% Sigma FBS
- 1% Pen/Strep
- 0.5 uM Human Insulin
- 0.1uM Dexamethasone
- 0.5 mM IBMX
- 33 uM Biotin
- 2 nM T3
- 30 uM Indomethacin
- 17 uM Pantothenate